

doi:10.1088/0264-9381/28/14/148001

M. Azreg-Aïnou, *Class. Quantum Grav.* **28** (2011) 148001 (2pp)

Comment on ‘Spinning loop black holes’

Mustapha Azreg-Aïnou

Başkent University, Department of Mathematics, Bağlıca Campus, Ankara, Turkey

Abstract

We review the derivations and conclusions made in Caravelli and Modesto (2010 *Class. Quantum Grav.* **27** 245022) and show that most of the analysis performed there is not valid.

PACS numbers: 04.60.Pp, 04.70.-s

In subsection 4.2 of [1], the authors, upon performing a type 2 complexification on a spherically symmetric metric in retarded Eddington-Finkelstein (EF) coordinates, arrived at

$$ds^2 = Gdu^2 + 2dudr - \Sigma d\theta - 2a \sin^2 \theta dr d\phi + [a^2(G - 2) \sin^2 \theta - \Sigma] \sin^2 \theta d\phi^2 + 2a(1 - G) \sin^2 \theta d\phi du, \quad (1)$$

where we have added the term $2dudr$ missing in Eq. (22) of [1]. G and Σ are functions of (r, θ)

$$\Sigma(r, \theta) = \rho^2 + \frac{a_0^2}{\rho^2}, \quad G(r, \theta) = \frac{\rho^2 - 2mr}{\Sigma} = \frac{\Delta - a^2 \sin^2 \theta}{\Sigma} \quad (2)$$

$$\text{with } \rho^2 = r^2 + a^2 \cos^2 \theta, \quad \Delta = r^2 - 2mr + a^2. \quad (3)$$

A claim was made in the first paragraph of page 14 of [1] that¹ metric (1) can be written in the Boyer-Lindquist (BL) coordinates (where the metric coefficients of the mixed terms $dt dr$ and $d\phi dr$ are zero) and be brought to Eq. (26) of [1], with Σ as given in (2), by a ‘coordinate transformation’ of the form

$$du = dt + g dr, \quad d\phi = d\phi' + h dr, \quad (4)$$

where g and h are supposed to depend only on r . We show that this is not possible and that all the analysis and conclusions made in [1], which are based on this claim, are not valid.

If we substitute (4) into (1), then the requirement that $g_{tr} = 0$ and $g_{\phi'r} = 0$ leads to

$$h(r) = -\frac{a}{\Sigma G + a^2 \sin^2 \theta} = -\frac{a}{\Delta(r)}, \quad g(r, \theta) = -\frac{\Sigma + a^2 \sin^2 \theta}{\Sigma G + a^2 \sin^2 \theta} = -\frac{r^2 + a^2}{\Delta} - \frac{a_0^2}{\Delta \rho^2}. \quad (5)$$

Since g depends on θ through ρ^2 , the system of equations (4) does not constitute a coordinate transformation. Thus, Eq. (26) of [1], with Σ as given in (2), does not describe the geometry of solution (1) in BL coordinates.

¹In that paragraph the expression ‘ $\Sigma(r, \theta)$ defined in (21)’ should read ‘ $\Sigma(r, \theta)$ defined in (44)’.

Figures 8 and 9 of [1] plotting the Ricci scalar, which have been sketched using Eq. (26) of [1] with Σ defined by (2), Eq. (45) and the last paragraph of page 14 of [1] are not valid. Figures 10, 11 and 12 of [1] are likely not valid too.

We have derived an expression for the Ricci scalar of the solution using metric (1)

$$R(r, \theta) = \frac{2a_0^4[\rho^4 + (r^2 + 2mr)\rho^2 - 6mr^3]}{\rho^2(\rho^4 + a_0^2)^3} + \frac{4a_0^2\rho^2[(4a^2 + 3r^2 - 3a^2\cos^2\theta)\rho^2 - 6mr^3]}{(\rho^4 + a_0^2)^3}, \quad (6)$$

and found it different from that plotted in figures 8 and 9 of [1]. In the limit $r \rightarrow 0$ and $\theta \rightarrow \pi/2$, we have $\lim_{(r,\theta) \rightarrow (0,\pi/2)} R(r, \theta) = 0$, which is not equal the limit given in the caption of figure 9 of [1]. A plot of the Ricci scalar (6), as shown in figure 1, is manifestly different from that of figure 9 of [1].




Figure 1: The Ricci scalar $R(r, \theta)$ of metric (1) for $m = 10$, $a = 5$ and $a_0 = 0.1$.

Similarly, the metric given in Eqs. (50) and (51) of [1] cannot be converted to BL form by transformation (4), since this would require a dependence on θ of both functions h and g . Thus, Eq. (55) of [1] and the claim made in the paragraph preceding it, as well as any conclusion based on Eq. (55) of [1], are not valid. Our conclusion extends most likely to Eq. (72) of [1], which has been derived using (4) as a ‘coordinate transformation’ when h and/or g depend(s) on θ .

References

- [1] Caravelli F and Modesto L 2010 *Class. Quantum Grav.* **27** 245022–31